

## Masks

Masks in healthcare workers;  
-Surgical masks and respirators (N95) appear similar in preventing viral infections, [N95 masks having slightly – not statistically different- lower infection rates (~1-2%).  
-Cloth masks are poorer than surgical (with ~2% RTI x4 wks).  
-No RCTs examined transmission to others or COVID-19.  
-Masks just one-part PPE and transmission precautions.

Mask in the community may reduce transmission of viral RTI (from 2 RCTs). If community risk was ~25% over 6 weeks, masks could decrease that to ~19%. No COVID-19 research, many studies examined others risk once someone was sick, and the overall certainty of evidence is low. Any mask use should be combined with social distancing and other preventive strategies

## Drug Treatment

Without further evidence, hydroxychloroquine is not appropriate for patients with COVID-19 in primary care. A number of recent trials/studies show an increased risk of side effects and QT prolongation especially at higher doses. RCTs are ongoing and hopefully they will provide more insight into the benefit/harm of this empiric treatment.

To date, no published RCTs have demonstrated benefit of treating COVID-19 patients with remdesivir, lopinavir–ritonavir or oseltamivir. One interim analysis of remdesivir suggests improved time to recovery. Full publication of studies and ongoing trials will help to answer this question.

## Clinical Factors

- Cough, fever and dyspnea are the most common symptoms of COVID-19.  
-At least 80% of cases are clinically mild, ~10% are hospitalized and 25% of admitted patients require intensive care.  
- Mortality risk factors include long-term care residents, age >65, co-morbid illnesses, and COVID-19 associated cardiac injury.

Transmission of COVID-19 can occur in people who are currently asymptomatic (including those who will remain asymptomatic and those who are early and not symptomatic yet). Case reports suggest this occurs in 6-13% of cases, although modelling suggests this might be higher. ~50% of carriers are asymptomatic when an entire population is tested.

## Testing

Studies of clinical PCR sensitivity are limited and vary widely for many reasons. Even if test sensitivity ranged between 50-90%, patients with low pretest probability (example 10%) would have at worst a 5% false negative rate.

While IGM and IGG antibodies (serology) may tell and individuals recent or past exposure – it is unclear whether antibodies confer immunity to subsequent infection. Accuracy of antibody testing likely requires validation in large number of infected and non-infected individuals.

## Miscellaneous

There is no reliable evidence that NSAIDs, ACE inhibitors or ARBs increase the risk of COVID-19 or affect disease severity/mortality from COVID-19

Unfortunately, no specific technique, including the Roth Score, reliably assures dyspneic patients are safe. No studies assessed dyspnea in COVID-19 patients. Clinicians are encouraged to use available tools (**BMJ Virtual Assessment tool**) and have patients assessed in-person if any concerns.